

- 1] For a finite non empty set of primes  $P$ , let  $m(P)$  denote the largest possible number of consecutive positive integers, each of which is divisible by at least one member of  $P$
- (i) Show that  $|P| \leq m(P)$ , with equality if and only if  $\min(P) > |P|$
- (ii) Show that  $m(P) < (|P| + 1)(2^{|P|} - 1)$
- (The number  $|P|$  is the size of set  $P$ )
- 2] For each positive integer  $n$ , find the largest integer  $C_n$  with the following property. Given any  $n$ -real valued functions  $f_1(x), f_2(x), \dots, f_n(x)$  defined on the closed interval  $0 \leq x \leq 1$ , one can find numbers  $x_1, x_2, \dots, x_n$ , such that  $0 \leq x_i \leq 1$  satisfying  $|f_1(x_1) + f_2(x_2) + \dots + f_n(x_n) - x_1 x_2 \dots x_n| \geq C_n$
- 3] Let  $A_1 A_2 A_3 A_4$  be a quadrilateral with no pair of parallel sides. For each  $i = 1, 2, 3, 4$ , define  $\omega_i$  to be the circle touching the quadrilateral externally, and which is tangent to the lines  $A_{i-1} A_i, A_i A_{i+1}$  and  $A_{i+1} A_{i+2}$  (indices are considered modulo 4 so  $A_0 = A_4, A_5 = A_1$  and  $A_6 = A_2$ ). Let  $T_i$  be the point of tangency of  $\omega_i$  with  $A_i A_{i+1}$ . Prove that the lines  $A_1 A_2, A_3 A_4$  and  $T_2 T_4$  are concurrent if and only if the lines  $A_2 A_3, A_4 A_1$  and  $T_1 T_3$  are concurrent.
- 4] Determine whether there exists a polynomial  $f(x_1, x_2)$  with two variables, with integer coefficients, and two points  $A = (a_1, a_2)$  and  $B = (b_1, b_2)$  in the plane, satisfying the following conditions.
- (i)  $A$  is an integer point (i.e  $a_1$  and  $a_2$  are integers);
- (ii)  $|a_1 - b_1| + |a_2 - b_2| = 2010$ ;
- (iii)  $f(n_1, n_2) > f(a_1, a_2)$  for all integer points  $(n_1, n_2)$  in the plane other than  $A$ ;
- (iv)  $f(x_1, x_2) > f(b_1, b_2)$  for all integer points  $(x_1, x_2)$  in the plane other than  $B$
- 5] Let  $n$  be a given positive integer. Say that a set  $K$  of points with integer coordinates in the plane is connected if for every pair of points  $R, S \in K$ , if there exists a positive integer  $l$  and a sequence  $R = T_0, T_1, T_2, \dots, T_l = S$  of points in  $K$ , where each  $T_i$  is distance 1 away from  $T_{i+1}$ . For such a set  $K$ , we define the set of vectors  $\Delta(K) = \{\overrightarrow{RS} | R, S \in K\}$ . What is the maximum value of  $|\Delta(K)|$  over all connected sets  $K$  of  $2n + 1$  points with integer coordinates in the plane?
- 6] Given a polynomial  $f(x)$  with rational coefficients, with degree  $d \geq 2$ , we define a sequence of sets  $f^0(\mathbb{Q}), f^1(\mathbb{Q}), \dots$  as  $f^0(\mathbb{Q}) = \mathbb{Q}, f^{n+1}(\mathbb{Q}) = f(f^n(\mathbb{Q}))$  for  $n \geq 0$ . (Given a set  $S$ , we write  $f(S)$  for the set  $\{f(x), x \in S\}$ )
- Let  $f^\omega(\mathbb{Q}) = \bigcap_{n=0}^{\infty} f^n(\mathbb{Q})$  be the set of numbers that are in all of the sets  $f^n(\mathbb{Q})$ . Prove that  $f^\omega(\mathbb{Q})$  is a finite set.